



KING HENRY VIII SCHOOL

Medical Policy and Procedures July 2022

Name of policy	Date reviewed	By whom	Next review	Responsibility
Medical	July 2022	S Cadwallader School Nurse Dr. M B Cuthbert Deputy Head	Annually July 2023	S Cadwallader

Contents

Policy for Supporting Pupils with Medical Needs	3
Policy for Administering Medicines.	4
Procedure for Dealing with First Aid Emergencies/Illness during School Hours	7
Medical Emergency Procedure including list of first aiders and first aid box contents	8
Information on Anaphylaxis and School Protocol.	11
Information on Asthma and School Protocol.	15
Information on Diabetes Mellitus and School Protocol.	18
Information on Epilepsy and School Protocol.	21
Protocol for concussion	23
Policy for children at risk of an Eating Disorder	26
Policy for children at risk of Self-Harm	30
Procedures for blood & body fluid spillage.	33
Reference list	34
<u>Appendix</u>	
Homely Remedy Protocol. (for paracetamol, Ibuprofen, Piriton, Throat Lozengers, bite cream)	35
Care Plan for pupils at risk of an Anaphylactic Reaction.	45
Instruction for using the EpiPen. (AAI)	48
Consent form for using Adrenaline Auto Injector	49
Consent form for emergency use of Salbutamol inhaler	50
Care Plan for Pupils at risk of an Asthma Attack.	51
Medicines in School Pro-forma	54
Annual Health Care pro-forma	55

POLICY FOR SUPPORTING PUPILS WITH MEDICAL NEEDS AT KING HENRY VIII SCHOOL.

1. Pupils with medical conditions will have the same rights of admission to King Henry VIII School as other children.
2. The school will ensure that pupils with medical needs will receive proper care and support whilst in the school environment.
3. Basic information and protocols will be available to all staff and parents for the most common medical conditions such as Asthma, Diabetes, Epilepsy and Severe Allergic Reactions (Anaphylaxis).
4. A protocol will be available for all other First Aid emergencies and should be read by all staff.
5. All members of staff will have access to a copy of the school handbook which will contain protocols and policies regarding medical conditions.
6. Details of children with common medical conditions will be available on iSams
7. It is the responsibility of all staff to read protocols and policies and notice board in the senior room common room regarding medical information.
8. All medical policies and protocols will be reviewed annually to ensure up to date information is available.
9. All Children with a Medical Condition will have an Individual Health Care Plan.
10. All staff will have access to the Individual Health Care Plans should a child with a medical condition be in their care. This will also be available on iSams
11. A copy of the Individual Health Care Plan must be taken on all educational visits should an emergency arise.
12. All individual health Care Plans will be reviewed annually to ensure up to date information is available.
13. The School Nurse is the first contact for parents and staff in regards to the medical needs of students.
14. The School Nurse will liaise with external agencies and will inform appropriate staff of the outcome should it be required.
15. Any medical intervention that takes place should be documented in the pupil's medical file. This is in the medical room and can be accessed via the school nurse to maintain confidentiality.

16. It is the parent/guardian responsibility to inform the school should there be any changes to their child's medical needs during the school year. This is to allow the school to update the medical files and provide the best support for their child.
17. Children with medical conditions should be allowed to participate in all physical activities and extra-curricular sport. There should be a flexible approach to allow students to partake in line with their own abilities. If there are any restrictions in relation to physical activity, then this will be noted in their Individual Health Care Plan.
18. All staff must treat medical information as confidential and must support children in their care promoting independence and maintaining dignity at all times.
19. Training will be available for all staff in regard to Anaphylaxis, Asthma, Epilepsy and Diabetes; this can be accessed via the School Nurse, and online <https://allergywise.org.uk/>
<https://learn.epilepsy.org.uk/> <https://www.virtual-college.co.uk/resources/free-courses/awareness-of-type-1-diabetes> <https://sch.educationforhealth.org/wp/elearning/>
20. It is the Parent / guardian responsibility to inform the school should be changes to health requiring the use of crutches, or inability to write. A temporary disabled care plan will be completed by the school nurse prior to return to school, to ascertain ability, needs, changes to school routine to accommodate

POLICY FOR ADMINISTERING MEDICINES AT KING HENRY VIII SCHOOL

1. Medication should not be carried around school. However, pupils with Asthma may carry their inhalers and for those at risk of an anaphylactic reaction may carry their adrenaline auto injector.
2. A child's medication should be given and stored in school if it is deemed detrimental for it to be omitted during the school day. (DfES 2005)
3. Consent - Pupils who require prescribed medication to be taken during the school day must bring it straight to the School Nurse at the start of the school day. A prescribed medication form must be completed by the parents, explaining dosage, time of administration, the last time medication was taken and whether the pupil is able to self-medicate. The form is available on the school website for parents to download for this purpose (See Appendix 5). All medication should be kept in its original container with the prescribing label still attached. All medication must be prescribed by a doctor with the exception of General Sales List (over the counter) medication. This may be sent into school with prior consent and agreement from the School Nurse ensuring the medication form is completed.
4. Should a pupil become ill or be injured during the school day they must be sent to the school nurse for assessment accompanied by another pupil. The School Nurse will decide whether the pupil is well enough to remain in school or if the best option is to contact parents to take home.
5. The nurse may administer medication under the Homely Remedy Protocol (see appendix 1) to allow for the child to remain in school with prior written consent from the parents. These will include Paracetamol, Ibuprofen and Piriton,(Antihistimine). Lozenges, anthisan, Burn Shield gel.
6. The school nurse may administer prescribed or General Sales List medication. In the absence of the nurse, only authorised personnel with appropriate training can administer medication (DfES 2005)
7. All medication will be locked in a secure cupboard only accessible by the School Nurse and authorised personnel. Emergency Inhalers and adrenaline auto injectors will be stored in an unlocked filing cabinet in the Medical Room named **Medication Held in School**. These can be accessed by all staff in accordance with (DfES 2005), for emergency use only.
8. For administering medication on school trips, see separate policy.
9. The School Nurse is responsible for the safe storage of medication and will keep a list with expiry dates. These will be updated regularly.
10. For medication held in school it is the responsibility of the parents to keep note of expiry dates and supply new medication when these expire.

11. All pupils with a chronic medical illness will have an Individual Health Care Plan. These can be found in the medical folder in the Medical Room.
12. It is the responsibility of the parent/guardian to inform the school should there be any changes to their health needs and any change in medication. This will allow for the correct support to be given in school.
13. Any medication that is given in school will be documented, noting Pupil name, date of birth, name of drug, dosage and time given.
14. In the case of medication being brought in to school, it is the person administering the drugs responsibility to check that the right drug is given to the correct person. This can be done by cross checking personal information such as name, date of birth, medication label and consent form.
15. If you are in any doubt about medication to be given then you must not give it until you have clarified your doubts. In the absence of the School Nurse you must contact the child's parents, with regards to medication.

Medications commonly used to treat ADHD.

Ritalin(methylphenidate), dexamphetamine (Dexedrine) Atomoxetine (Strattera) Concerta, Tranquilyn, equasym

Licensed for children 6 yrs. and over currently.

ADHD is a complex phenomenon involving many contributing factors. Therefore, assessment and diagnosis is usually made by a specialist child psychiatrist or paediatrician. The symptoms can involve extreme persistent behaviour, such as over-activity, restlessness, impulsiveness, and inability to concentrate. The behaviour is apparent in more than one setting, for example home and at school.

The medication works by stimulating parts of the brain responsible for consciousness and control of attentions and activity, thus increasing concentration ability and decreasing restlessness in children who are overactive, impulsive and easily distracted.

A child who has been prescribed a controlled drug may have it in their possession but due to its potential for misuse it is advisable that it is stored securely so that the safety of other pupils is not put at risk. These medications are Class B controlled drugs and under the requirements of Misuse of Drugs (safe control) Regulation as of 1971 amendments 2010 should be stored in a double locked cabinet within a designated area where staff with authorised access should hold keys.

It is therefore vital that any Medication brought into the school be appropriately managed and stored. Accurate records of the dispensing to the medication should be made.

To assist in administering at school Parents should supply letters from the Consultant outlining

Name
Dosage
Frequency
Diagnosis
Review date

Parents must also complete a Prescribed Medication Form (should be available for Parents to download from the website. Any changes to doses should be passed on through this route accompanied by a letter from Consultant.

Receiving Ritalin (concerta, equasym, Tranquilym) for storage in School

Medicines should be in their original packaging and be clearly marked with the child's name and prescriber instructions.

A designated member of staff (teacher, learning assistant, office staff, Nurse) should record the amount received, the name of the child for whom it is intended, the expiry date and prescriber instructions.

The designated member of staff and the Child's parent or carer should both sign to confirm medicine had been handed over to the school. Expired / Unused medicine should be returned to the parent or carer as a matter of routine, whether weekly, monthly or at end of half term. Both parent / carer and staff member should sign to say this has been done.

Side Effects

The main side effects are reduced appetite and staying awake late.

Other less common side effects are anxiety, headaches, nervousness, headache, stomach ache, dizziness

Administration of Ritalin

The member of staff should always check that the child's name and the dose of medication prescribed match what is written on the container and support plan.

The member of staff should supervise the self-administration of the medicine at a time and place agreed with pupil, parent and other staff member. Staff should ensure medicine has been taken.

If a child refuses the medication a note should be made in the record and parents informed.

The member of staff should record the amount of medication taken and the time it was taken.

Best practice is that a second member of staff (or pupil) should countersign the entry in the medication record book. Both staff / pupil should check the remaining amount is correct and accurately recorded.

All controlled drugs must be checked, administered, and signed for by two persons. Stock levels and details of administration are recorded in the Controlled drug book as well as pupil held documents.

PROCEDURE FOR DEALING WITH FIRST AID EMERGENCIES/ILLNESS DURING SCHOOL HOURS AT KING HENRY VIII SCHOOL.

Illness

- Any pupil taken ill during the school day must be accompanied by another pupil or member of staff and taken to the medical room. In the absence of the school nurse, please report to reception, they will locate the school nurse or First Aider.
- The nurse will inform the appropriate members of staff of any action taken.
- Under no circumstances should the pupil contact their parents directly
- Should it be decided that it is better for the pupil to return home, then the school nurse or reception staff will make contact with the parents/guardian.
- Should the child be sent home due to vomiting or diarrhoea then the parents/guardians must be informed not to return their child to school for 48 hours following the last episode. (In line with Public Health England Guidelines)

Accidents

- Minor accidents or injury should be dealt with as for illness.
- In the event of a more serious accident/injury, do not attempt to move the casualty but contact the School Nurse on **199** or **07483 065527** or follow the procedure for medical emergencies if deemed necessary.
- In the absence of the School Nurse, contact a qualified First Aider, a named list with contact numbers is available in all departments, reception, senior common room, and medical room.
- If it is deemed appropriate an ambulance should be called by **any** member of staff, following the Medical Emergency Procedure.
- Parents/guardians/next of kin will be contacted and advised to attend the named accident and emergency department.
- A responsible member of staff must accompany the casualty if less than 16 years of age with the ambulance crew to hospital and remain until a parent/guardian arrives.

An accident report should be completed for all accidents and near misses by the member of staff that witnessed the incident. All the required information must be completed including personal details and then passed to the Health and Safety Officer who will decide whether it is reportable to RIDDOR. Accident forms can be located in the Medical Room or the Senior Common Room. On no account should completed forms be left in the accident book, this is for confidentiality reasons in line with the Data Protection Act 1998 and reporting criteria.

When to call an Ambulance

Once you have completed your first aid assessment and you identify the need; or if at any time you feel unable to administer First Aid confidently to a pupil a member of staff or visitor; or you have not received the appropriate training then an ambulance should be called using the following procedures.

MEDICAL EMERGENCY PROCEDURE FOR KING HENRY VIII SCHOOL

☒ Dial 999 (if using an internal phone dial 9 for outside line first) ☒ Ask for the appropriate service: Ambulance, Fire Brigade or Police. ☒ Make sure you have the following information available:

- name
- The incident information that has occurred
- Your telephone number – 0247 627 1111
- The location of the incident.

For Main School Site.

King Henry VIII School
Warwick Road
CV3 6AQ

For Stonebridge Highway

King Henry VIII Playing Fields
Stonebridge Highway
Coventry
CV3 4EJ

Bablake Playing Fields / Pavillion
Hollyfast Road, Coventry CV6 2AE

For Music Block, Science Block, DT or School Playing Field

King Henry VIII School
Spencer Road
CV3 6AQ

For Foundation Astro

Astro Pavilion (Bablake Old Boys Pavilion)
Norman Place Road
Coventry
CV6 2ND

Astroturf Bablake
Dunscroft Avenue, Coventry, CV6 2BX

- Arrange for someone to meet the ambulance and direct to the appropriate location.
- A responsible member of staff should accompany the pupil to the hospital with the ambulance crew and remain with the pupil until the parent/guardian arrives.
- On no account should a member of staff attempt to take the pupil to hospital in their own vehicle.
- **Always contact the school nurse if you have to make an emergency call. It is now a requirement that these are logged by a phone call to the Health & Safety Executive.**

Body Fluids

All bodily fluids should be considered as potentially hazardous. General body fluids include fluid from cuts, scabs and skin lesions, blood, urine, faeces, vomit, nasal discharge, and saliva. Gloves and aprons should be always worn when there is a risk of encountering bodily fluids. Once used, gloves and aprons should be discarded in a bio hazardous bag, available from the biology prep

room. For spillages see procedure for dealing with Body fluids and Bloods Spillages. Covid PPE is available.

Correct Procedure for Removal of Gloves

- With right hand, pinch palm of gloves on left hand and pull left glove down and off fingers. Form left glove into a ball and hold securely in fist of right hand.
- Insert two fingers of left ungloved hand under inside rim of right glove on palm side, pull glove slowly down over fingers and over balled left glove.
- This can now be discarded in bio hazardous bag with any soiled dressings. ☑ Wash Hands

Contents of First Aid Kits at King Henry VIII School

- 1 Guidance Leaflet
- 20 Individually wrapped plasters
- 2 Sterile eye pads
- 4 Triangular bandages
- 6 Safety pins
- 2 Large unmedicated dressings
- 6 Medium unmedicated dressings
- 1 roll of micropore tape
- 1 pair disposable gloves
- Scissors
- Cleansing wipes
- 1 Ice pack
- 1 resuscitation face shield

Location of First Aid Kits at King Henry VIII School

- Medical Room
- Staff Room
- Chemistry Prep Room
- Sports Hall Office (Boys)
- Sports Hall Office (Girls)
- Biology Prep Room
- Main Hall Foyer
- Caretakers Room
- Library
- Art Department
- Physics Office
- Music Office
- Classics Office
- History Office
- IT4

- DT3
- DT2
- DT1
- Maintenance Office
- Sports Centre Office
- Sixth Form Centre

INFORMATION ON ANAPHYLAXIS AND SCHOOL PROTOCOL MANAGING SEVERELY ALLERGIC PUPILS IN SCHOOL.

Definition of Anaphylaxis

Anaphylaxis involves one or both of two features: -

- Respiratory difficulty (swelling of the airway or asthma).
- Hypotension (fainting, collapse, or unconsciousness).

(The Anaphylaxis campaign, 2009).

What's happening to the Body?

- An anaphylaxis reaction is caused by the sudden release of chemical substances, including histamine, from cells in the blood and tissues where they are stored.
- The release is triggered by the reaction between allergic antibodies and the allergen
- The person would have been exposed to the allergen (the thing that they are allergic to) previously. At that time, the body misjudged the allergen as a threat and started to make antibodies against it. The next time the body is exposed it over reacts and causes the above symptoms.
- Anaphylaxis is a severe systemic allergic reaction.
- At the extreme end of the allergic spectrum.
- The whole body is usually affected within minutes of encountering the allergen.
- It can take seconds or several hours for a reaction to occur.

(The Anaphylaxis campaign, 2009)

Allergic reactions can produce many unpleasant symptoms; only a few are likely to be described as anaphylaxis.

(Ewan 1998)

The Symptoms

- Swelling of the mouth or throat.
- Difficulty in swallowing or speaking.
- Alterations in the heart rate.
- Hives anywhere on the body.
- Abdominal cramps and nausea.
- Sudden feeling of weakness.
- Difficulty in breathing.
- Collapse and unconsciousness.

(The Anaphylaxis Campaign, 2009)

Types of Reaction

- Uni-phasic, meaning one phase. Develops rapidly, usually involving the airway or circulation. Once treated the symptoms go away and don't return.
- Bi-phasic, meaning two phases. 6% of children have bi-phasic reactions. It develops rapidly, is treated, and then there appears to be a rest period when all symptoms appear to have gone away for 1-2 hours. Breathing and circulation symptoms return and can become very serious.

(Lee and Greenes, 2000. The Anaphylaxis Campaign, 2009)

Common Causes

- Peanuts ☒ Wasp
- Tree nut ☒ Bee
- Milk ☒ Latex
- Egg ☒ Penicillin
- Sesame ☒ Blood Products
- Fish ☒ Some medication
- Shellfish ☒ Kiwi

Treatments

Adrenaline is the main treatment and does the following:-

- ☒ Reverses swelling
- Relieves Asthma
- Constricts the Blood Vessels
- Stimulates the heartbeats

Adrenaline works in seconds, whereas antihistamines take about 15 minutes. This is useful if the reaction is coming on slowly, asthma inhalers will also help at this time.

Incidentally, children with asthma as well as severe allergies are far more at risk of a severe reaction than allergic children who do not have asthma. (Sampson et al 1992)

Devices

Adrenaline auto-injectors: -

- Adult dose 0.3mg
- Child dose 0.15mgs
- Up to 2 years shelf life **Storage**
- Accessible
- Avoid extremes of temperatures; the devices are designed to be stored at room temperature. They remain stable up to about 40 degrees C. Should not be stored in the fridge or left in sunlight.
- Clearly labelled
- In date

Emergency Adrenaline Auto-Injector (AAIs) in School

Following guidance from the Department of Health – amendments to the Human Medicines Regulations 2017: Adrenaline-auto injectors, all schools are now able to obtain Adrenaline Auto injectors (AAIs) for emergency use without prescription. This is a discretionary power enabling

schools to do this if they wish. **Currently we do not hold an Emergency Adrenaline Auto- Injector due to issues of availability.**

- The Emergency Adrenaline auto injector will be held in the medical room and used under the School Nurse 's direction.
- The Emergency Adrenaline auto injector (AAI) is only available for Pupils known to be at risk of anaphylaxis, where both parental consent and medical authorisation has been given.
- The Adrenaline Auto injector (AAI) should be considered a spare/ back up device only and not a replacement for a pupils own AAI. (for example if their own has been used / broken.)
☒ In the event of possible severe allergic reaction in a pupil who does not meet the above Criteria, emergency services (9) 999 should be contacted and advice sought from them as to whether administration of the Spare emergency AAI is appropriate.
- Written parental consent for the use of the Emergency Adrenalin auto injector (AAI) needs to be obtained. A list of children with parental/ guardian consent will be kept with the emergency adrenaline auto injector (AAI)
- Use of the emergency adrenaline auto injector will be documented in the pupil's medical records.

The emergency Anaphylaxis Kit

- One or more AAI(s)
- Instructions of how to use the devices(s)
- Instructions on storage of the AAI device
- Manufacturers information
- A checklist of injectors identified by batch numbers and expiry date with monthly checks recorded
- A list of pupils to whom AAI can be administered
- An administration record

Allergen avoidance

- Ensure you know the child in your care and familiarise yourself with their allergies. The child and their parents will have become experts at this and will know what they need to avoid, so speak to the child or the School Nurse.
- These children are very normal and it is important that they join in as many activities as possible.
- Risk assessments should be carried out in regards to location and activities that the child is involved in, particularly trips away, cookery, science experiments and mealtimes.
- Special occasions such as Christmas, Easter and fund raising cakes sales can pose an increased risk to allergy sufferers due to different food being brought in to school. Please be vigilant and risk assess where needed.
- Plan ahead in regards to cookery lessons and science experiments and inform the child or parent to discuss their risks.
- It is good practice not to use Food Technology rooms as form rooms for the allergic pupils.
- Be aware of any empty boxes being brought into school for Art/technology etc they may have contained things such as crunchy nut cornflakes etc.

Health Care Plans (see appendix)

- It is the responsibility of the School Nurse that all pupils will have a health care plan, reviewed annually.
- The school nurse will give a copy to the pupil/parent/guardian, the tutor and one will remain in the Medical File in the medical room.
- The tutor's copy must remain in their register for the purpose of someone else registering their class.
- A named picture of all children who have adrenaline auto injectors will be displayed in the Senior Common Room.
- All staff including those on supply should be encouraged to check iSams regularly and familiarise themselves with these Pupils so that they will be prepared in the case of an emergency.

What to do in the case of an allergic reaction?

- Contact School Nurse immediately on 199 or 07483 065527, stating name, form and location of the pupil, the allergic reaction.

In the absence of the School Nurse

- Ensure a responsible person always stays with the pupil.
- Ask a responsible person to collect the emergency drug pack for the pupil. This is found in the medical room, ask a member of the reception staff to direct you to its location.
- **For mild reactions**, to include rash/hives, swelling of the lips, itching, and stinging sensation of the mouth administer prescribed antihistamine. Monitor pupil and contact parents to inform them of the event.
- **Should the pupil have a severe reaction** to include swelling inside the throat and mouth, faintness, loss of consciousness, intense anxiety, wheezing similar to asthma attack, abdominal cramps, nausea and vomiting and widespread hives. A responsible person should call for an ambulance (see Medical Emergency Procedure, 2009). Name and age of child should be given along with reason of call 'child having a severe anaphylaxis reaction'
- A second trained member of staff will administer the adrenaline auto injector to the pupil as prescribed by the doctor. (See appendix 3 for guidance)
- The date and time the drug is given should be documented (see policy for administering medicines in school, 2009) and the used pens kept in a container ready for the ambulance crew.
- Place the pupil in a position of comfort, always ensuring maintenance of airway.
- An accident report should be completed and forwarded to the Health and Safety Officer who will then report it to RIDDOR.

If you would like more information please contact the School Nurse or the Anaphylaxis Campaign
Tel. 01252 542029 or www.anaphylaxis.org.uk

INFORMATION ON ASTHMA AND SCHOOL PROTOCOL.
MANAGING ASTHMATIC PUPILS IN SCHOOL

Definition of Asthma

- Difficulty in breathing, with a very prolonged breathing out stage.
(Cleaver et al, 2006)

What's happening to the body?

- Asthma is a condition that affects the airways – the small tubes that carry air in and out of the lungs.
- When an asthmatic encounter something that irritates their airways, it causes unwanted symptoms (also known as an asthma trigger).
- It can take seconds or several hours for a reaction to happen once exposed to the asthma trigger.
(Asthma UK, 2009)
- The muscle that is found in the walls of the airway tightens or goes into spasm.
- The airways become narrow and the lining of them becomes inflamed and start to swell.
- Sometimes sticky mucus and phlegm builds up which can narrow the airways further.
(Cleaver et al, 2006)

The Symptoms

- Wheezing as the casualty breathes out.
- Difficulty in speaking and whispering
- Features of hypoxia (lack of oxygen) such as grey blue tinge to the lips, earlobes and nail beds (cyanosis)
- Appears distressed and anxious.
- Dry, tickly cough
(Asthma UK, 2009. Cleaver et al, 2006. Newman et al, 2001)

Asthma Triggers

- | | |
|------------------------------|--------------------|
| ☒ Animals | ☒ House-dust mites |
| ☒ Air pollutants | ☒ Medicines |
| ☒ Colds and viral infections | ☒ Moulds and fungi |
| ☒ Emotions | ☒ Pollen |
| ☒ Exercise | ☒ Smoking |
| ☒ Food | ☒ Weather |
| ☒ Hormones | |

What causes Asthma?

It is difficult to say what causes asthma exactly but what is known is that:

- If you have a family history of asthma, eczema, or allergies you are more likely to develop it.
- Family history combined with specific environmental factors can influence whether you get asthma.
- Modern lifestyle changes such as housing, diet and a more hygienic environment may have contributed to the rise in asthma in the last few decades.

- Research shows that smoking through pregnancy significantly increases the risk of a child developing asthma.
- Children whose parents smoke have an increased risk of developing asthma.
- Environmental pollution can worsen asthma symptoms and may play an important part in causing some asthma's.
- Adult onset of asthma can develop following a viral infection.
- Irritants in the work place can lead to a person developing asthma (occupational asthma).

Medication

- Reliever Inhalers (usually blue), taken to relieve asthma symptoms immediately. (bronchodilators)
 - Salbutamol is the most popular medication and can have various trade names such as, Ventolin and salamol. 100mcg per puff and 2 are usually taken.
 - Terbutaline can also be used, also known as Bricanyl. 500mcg per puff and 1 is usually taken.
 - Preventer inhalers (usually brown, burgundy, purple or orange).
 - Various medications called corticosteroids, such as, becotide, beclomethasone, flixotide, and seretide. These are of no use to relieve asthma symptoms once they have manifested, but are used to prevent symptoms occurring.
- (British National Formulary, March 2008)
- There are a few other medications that can be used to manage Asthma, but these are the most common.
 - All Asthmatics will be encouraged to carry their own medication around with them for use in an emergency. A second inhaler **may** be stored with the school nurse. See Policy for Administering Medicines in School, 2009.

Storage

- Accessible
- Avoid extremes of temperatures; the inhalers are designed to be stored at room temperatures. Should not be stored in the fridge or direct sunlight, this can prevent the inhaler from working correctly.
- Clearly labelled
- In date

Asthma Trigger Avoidance

- Ensure you are aware of any children in your care that have Asthma and familiarise yourself with their triggers. The child and their parents will be experts on this, so makes sure you speak to them if you are concerned.
- These children are very normal and it is important that they join in as many activities as possible.
- Risk assessments should be carried out in regards to location and activities that the child is involved in, particularly trips away, physical education, and cookery and science experiments.

Health Care Plans (see appendix4)

- It is the responsibility and desire of the School Nurse that all pupils will have an individual health care plan. (This will be discussed with the parents.) On some occasions it may be felt that it is not necessary by the parents. In this case school protocol on how to manage an asthma attack should be adhered to. Individual plans will be reviewed annually.

- The School Nurse will give a copy to the pupil/parent/guardian, the tutor and one will remain in the medical file in the medical room. .
- A list of all asthmatics will be displayed in the Senior Common Room.
- All staff including those on supply should be encouraged to check iSams regularly and familiarise themselves with these pupils so that they will be prepared in the case of an emergency.

Games/PE lessons and Asthma Management

- Ensure you know who has Asthma in your group and adhere to asthma trigger avoidance. ☒ Allow the pupil to increase their fitness levels gradually.
- It is the pupil's responsibility to have their inhaler with them when they undertake any physical activity.
- If exercise triggers the pupil's asthma then they should be reminded to take their inhaler immediately before they participate.
- Should the pupil complain of asthma symptoms during the activity, they must be allowed to stop immediately and be encouraged to take their reliever inhaler and wait until they feel better before they continue.
- When asthma is under control, pupils should be able to take part in all sports.
- Should a pupil be unable to participate in physical education regularly, then the School Nurse will contact the parents to offer advice on asthma management. The nurse will liaise with outside agencies such as Asthma UK, the pupils GP, asthma nurse or Consultant.

Emergency Salbutamol Inhaler in school

Following the guidance from the Department of Health, all schools are now able to obtain salbutamol inhalers for emergency use without prescription. This is a discretionary power enabling schools to do this if they wish.

- The emergency inhaler will be held in the medical room and only used under the school nurse's direction. It will be available to any pupil with asthma, or who has been prescribed an inhaler as reliever medication, and can be used if the pupil's prescribed inhaler is not available (for example, because it is broken, or empty).
- Written parental consent for use of the emergency inhaler needs to be obtained. A list of children with parental consent will be kept with the emergency inhaler. Disposable spacers will be stocked as well.
- If it is necessary for a child to use the emergency salbutamol inhaler, the information will be documented in the child's medical records.
- The emergency inhaler can be reused. The canister must be removed and the casing can be washed in warm soapy water and left to air dry before reconstructing.

What to do in the case of an asthma attack?

- Contact the School Nurse immediately on **199** or **07483 065527** **A responsible person will stay with the pupil and encourage them to:** ☒ Take 2 puffs of their reliever inhaler ☒ Sit up and loosen tight clothing.
- If no immediate improvement, continue to take 2 puffs (one at a time) every two minutes. Up to ten puffs can be taken.

In the event that the symptoms do not improve, a second responsible person should at this point call for an ambulance (see Medical Emergency Procedure, 2009) stating:-

- Reason of call – ‘child having a severe asthma attack’. ☑ Medication child has already received.

Then:

- Contact parents and tell them to go straight to the named hospital
- Encourage the pupil to continue to take 1 puff of their reliever inhaler every minute until Symptoms improve or the ambulance arrives.
- Ensure a responsible member of staff escorts the pupil with the ambulance crew to the hospital.
- An accident report must be completed and forwarded to the health and safety officer who will need to report it to RIDDOR.

If you would like more information please contact the School Nurse or Asthma UK, Tel. 0800 121 6244 or www.asthma.org.uk

INFORMATION ON DIABETES MELLITUS AND SCHOOL PROTOCOL MANAGING DIABETIC PUPILS IN SCHOOL

Definition of Diabetes Mellitus

- Diabetes Mellitus often referred to as diabetes, is a syndrome of disordered metabolism, usually due to a combination of hereditary and environmental causes resulting in abnormally high blood sugar. (Tierney et al 2002)

What is happening to the body?

- Glucose is the fuel used by the body for every cell. When not enough insulin is available or not able to function correctly, then the glucose cannot get into the cells where it is needed. This then builds up in the blood.
- The unused blood sugar circulates through the kidneys, when the amount is more than the kidneys can tolerate the extra glucose spills out into the urine. (Ministry of Health and Family Welfare Government of India, 2008-09)
- Glucose come from the digestion of foods containing carbohydrate such as bread, potatoes, chapattis, fruit, dairy products and other sweet foods. It is also produced by the liver.
- Without insulin, sugar accumulates in the blood and can cause hyperglycaemia.
- Insulin is vital for life and is a hormone produced by the pancreas, it helps the glucose to enter our cells where it is used as fuel for energy so we can work, play and generally live our lives.
- Diabetics must carefully balance the amount of sugar in their diet and regulate their blood sugar with insulin injections, tablets or diet; too much insulin or too little sugar can cause hypoglycaemia.

(Diabetes UK, 2015)

Undetected or unmanaged diabetes can lead to organ failure, coma and even death.

The Symptoms

- **Hyperglycaemia** o Dry skin. o Rapid Pulse.
 - o Deep, laboured breathing. o A smell of acetone or pear drops on the casualty's breath. o Excessive thirst. o Passing large amounts of urine.
 - o Tiredness and irritability.
- **Hypoglycaemia** o Hunger o Sweating o Drowsiness
 - o Pallor (pale) o Shaking
 - o Lack of concentration
 - o Irritability o Lack of responsiveness reduces
 - o May become unconscious.
 - o Cold clammy skin o Shallow Breathing.

(Diabetes UK, 2015, Cleaver et al, 2006, Newman et al, 2001)

Diabetes Types

- **Type 1 Diabetes.** o This develops if the body is unable to produce any insulin. o Usually appear before the age of 40. o The least common of the two types, accounts for about 5-15% of all people with diabetes.
 - o Unpreventable.

- **Type 2 Diabetes.** o Develops when the body can still make some insulin or when the insulin that is produced does not work properly.
 - o Most cases are linked to being overweight. o Usually appears after the age of 40. (in South Asian and Afro Caribbean people often appears after the age of 25) o Recently, more children have been diagnosed some as young as 7.
 - o The more common of the two types, accounts for about 85-95% of all people with diabetes.
 - o Some cases can be preventable.

(Diabetes UK, 2015)

Treatments

- Insulin pumps that infuse insulin 24 hours and are attached to the pupil. ☒ Insulin injections.
- Tablets
- Diet controlled.
- All are usually managed independently.
- Glucose tablets
- Dexrologel

Storage and disposal

- Insulin is usually stored in the refrigerator but can vary depending on the type of insulin. Advice is to remove 1 hour prior to administration.
- Tablets should be stored at room temperature.
- Any needles that are used should be disposed of in the yellow sharps box that is located in the Medical Room.

Awareness of Hypoglycaemia and management.

- Ensure you are aware of any children in your care that are diabetic.
- These children are very normal and it is important that they join in as many school activities as possible.
- Risk assessments should be carried out in regards to location and activities that the child is involved in, particularly trips away and physical education.

Health Care Plans

- It is the responsibility of the School Nurse to ensure that all pupils will have an Individual Health Care Plan.
- The School Nurse will give a copy to the pupil/parents/guardian, and one will remain in the medical file in the medical room.
- A list of all diabetics will be displayed in the Senior Common Room.

- All staff including those on supply should be encouraged to check the board regularly and familiarise themselves with these pupils so that they will be prepared in the case of an emergency.
- Pupils with diabetes will be encouraged to carry an emergency pack containing carbohydrate snacks, glucose tablets or dextroglucose should their blood sugar become low.
- The nurse will have a supply of glucose tablets in the medical room in the case of an emergency.

What to do should a pupil have a hyper/hypoglycaemic episode.

- Hyperglycaemia is unlikely to cause any problems whilst in school but if any of the symptoms are noticed the School Nurse must be informed immediately.
- If a Hypoglycaemic episode is suspected you should contact the school nurse immediately on **199** or **07483 065527**.
- The pupil should be encouraged to eat their emergency snack or have 3-5 glucose tablets or dextroglucose if able to swallow.
- Once the pupil recovers, parents will be contacted for further instruction.
- ❓ **Should the pupil deteriorate despite using their emergency pack or becomes unconscious at any point then an ambulance should be called. (See Medical Emergency Procedure, 2009) stating:-**
 - Diabetic pupil having a hypoglycaemic attack.
 - Emergency pack used. **Then**
 - Contact parents and tell them to go to the named hospital.
 - Ensure a responsible member of staff escorts the pupil with the ambulance crew to the hospital.
 - An accident report must be completed and forwarded to the health and safety officer who will need to report it to RIDDOR.

If you would like more information please contact the School Nurse or Diabetes UK, Tel. 020 7424 1000 or www.diabetes.org.uk

INFORMATION ON EPILEPSY AND SCHOOL PROTOCOL MANAGING EPILEPSY IN SCHOOL.

Definition of Epilepsy.

- A tendency to have repeated seizures or fits that start in the brain. It is a neurological condition as it affects the brain, but also a physical condition as it affects the body. (The National Society for Epilepsy, 2009)

What is happening to the Body?

- The cells in the brain, known as neurons, communicate with each other by using electrical impulses. During a seizure these are disrupted, which can cause the brain and the body to behave strangely.
- The severity of the seizures can differ from person to person. Some people will experience a trance like state for seconds or minutes whereas others will lose consciousness and have convulsions (uncontrollable shaking of the body).

(NHS.UK, 2009)

- The brain is responsible for all the functions of the body. What is experienced during a seizure will depend on where in the brain the epileptic activity begins and how widely and rapidly it spreads. Epileptic seizures will be unique to the individual.

(Epilepsy Action, 2009)

Types of Seizures and Symptoms

Seizures are divided into two main groups as follows:- Generalised seizures.

- Tonic clonic seizure - most common type of generalised seizure, consciousness is lost, the whole body stiffens and shakes (convulses). This is due to uncontrollable muscle contractions.
- Absence seizures – Brief loss of consciousness or awareness. No convulsions or falling over and lasts only seconds, usually seen in children.
- Myoclonic seizure – Sudden contractions of the muscles which cause a jerk. These can affect the whole body but usually occur in just one or both arms.
- Tonic seizure – Brief loss of consciousness, may become stiff and fall to the ground.
- Atonic seizure – Become limp and collapse, often with only a brief loss of consciousness.

Partial Seizures.

- Simple partial seizures – May have muscular jerks of strange sensations in one arm or leg. Can get an odd taste; develop pins and needles in one part of the body. Do not lose consciousness.
- Complex partial seizures – May behave strange for a few seconds or minutes, e.g. may fiddle with an object, or mumble, or wander aimlessly. As well as odd emotions, fears, feelings, visions or sensations. Consciousness is affected and you may not remember having a seizure.

Cleaver et al, 2006. Patient UK, 2009)

Common Triggers

- Tiredness, Lack of Sleep.
- Stress.
- Alcohol.
- Noncompliance with medication.
- Less common, flashing or flickering lights or patterns, affects only 5% of epileptics and is called photosensitive epilepsy.

Treatments

- Anti-epileptic drugs – aim to prevent seizures from happening but do not cure it.
- Rescue medication such as diazepam, administered during a seizure. ☒ Other options include vagus nerve stimulation.
- Surgery.

Storage

- Accessible
- Avoid extremes of temperatures. Stored at room temperature.
- Clearly labelled.
- In date.

Awareness and management of Epilepsy.

- Ensure you know the children in your care that have epilepsy and familiarise yourself with their triggers.
- These children are very normal and it is important that they join in as many activities as possible.
- Risk assessments should be carried out in regards to location and activities that the child is involved in, particularly Science, Design and Technology, Food Technology and trips away from school. **Health Care Plans.**
- It is the responsibility of the School Nurse that all pupils will have a health care plan.
- The School Nurse will give a copy to the pupil/parents/guardian, the tutor and one will remain in the medical file in the medical room.
- The tutors copy must remain in their register for the purpose of someone else registering their class.
- A list of all epileptics will be displayed in the Senior Common Room.
- All staff including those on supply should be encouraged to check the board regularly and familiarise themselves with these pupils so that they will be prepared in the case of an emergency.

What to do in the case of a convulsive seizure.

- Contact the school nurse immediately on **199** or **07483 065527**, stating name, form and location of the pupil.
- Protect the person from injury, i.e. remove all harmful objects from nearby.
- Place a cushion or coat underneath the pupils head for protection.
- Do not try to move the pupil unless in immediate danger.
- Do not try to restrain the pupil.
- Make sure pupils are not crowding around and remove them from the area if possible.
- Whilst convulsing the pupil may become incontinent, ensure you are discreet and maintain their dignity.
- Do not put anything in the pupil's mouth or give anything to eat or drink until fully recovered.
- Place in the recovery position once the seizure has finished. Reassure the pupil as they may not know what has happened and contact the parents informing them of the event

Call for an ambulance if:-

- You know it's the person's first seizure.
- The seizure continues for more than 5 minutes.
- A tonic-clonic seizure follows another without the person regaining consciousness.

- The person is injured during the seizure.
- You believe the pupil needs urgent medical attention.
- If an ambulance is called (See Medical Emergency Procedure, 2009). State the pupils name, age and reason of call – ‘child having an epileptic seizure’.
- Contact the parents and tell them to go to the named Accident and Emergency Department.
- A responsible member of staff should escort the pupil with the ambulance crew to the hospital.
- An accident report should be completed and forwarded to the Health and Safety Officer who will then report it to RIDDOR.

For all other seizures contact the School Nurse. In the absence of the school nurse the parents must always be contacted to inform them of the event. Should the pupil feel unwell the parents will take them home to recover.

If you would like more information please contact the School Nurse or The National Society for Epilepsy. Tel. 01494 601 400 or www.epilepsy.or.uk

INFORMATION ON CONCUSSION IN SPORT & SCHOOL PROTOCOL

What is concussion?

A concussion is a disturbance in brain function caused by a direct blow to the head or indirect force to the body, induced by biomechanical forces. The medical term for concussion is minor traumatic brain injury. Loss of consciousness occurs in than less than 10% of concussions, it is not a requirement for diagnosing concussion. Standard brain scans are typically normal with someone with concussion.

Signs and symptoms of concussion

Symptoms of concussion may be delayed and only present 24-48 hours after the injury.

The most common symptoms of concussion are:

- Headache
- Dizziness
- Nausea
- Loss of balance
- Confusion, such as being unaware of your surroundings
- Feeling dazed with a “foggy head”.
- Disturbances with vision, such as double vision or seeing 'stars' or flashing lights ☒
- Difficulties with memory

Less common signs and symptoms

- Loss of consciousness
- Vomiting
- Slurred speech
- Vacant or blank stare
- Changes in behaviour, such as feeling unusually irritable
- Inappropriate emotional responses, such as suddenly bursting into laughter or tears

Visible clues may also indicate a concussion

- Lying motionless on the ground or slow to get up
- Unsteady on their feet
- Grabbing or clutching their head

Signs of concern

- A neck injury is suspected or a player complains of severe neck pain
- Pins & needles in arms or hands
- Convulsion or seizure
- Deteriorating conscious state

If any player has any of the above symptoms, an ambulance must be called. Neck pain or pins & needles may indicate a cervical spinal fracture and the player must be removed from the field of play by emergency healthcare professionals with appropriate spinal care training.

Who is most at risk?

Children and adolescents are more susceptible to concussion due to the continuing development of the brain and should be treated more conservatively. Someone with a history of two or more concussions within the past year are at greater risk of further brain injury and slower recovery.

There are several factors that cause vulnerability in a head injury

- Previous concussion. Repeated concussions can cause cumulative brain damage leading to second-impact syndrome
- Alcohol or drug use
- Having a condition that makes you bleed more easily – haemophilia
- Taking anticoagulant medication such as warfarin

High risk groups

- Young men
- Sportspeople
- Those with a history of mental health problems

It is important to be aware of demographic groups at particular risk of head injuries and to consider whether symptoms are related to an undiagnosed, previously unreported injury

Management of concussion

Anybody who has received a heavy jolt to the body or head, or a direct blow to head should be removed from play.

Advised to seek further medical intervention, Off Games for minimum 2 Weeks or Dr's not advising fit to Play.

Questions to ask a suspected concussed player Failure to answer may suggest a concussion

- ✓ What venue are we at today?
- ✓ Which half is it now?
- ✓ Who scored last in this game?
- ✓ What team did you play last week? ☐ Did your team win the last game?

Parents need to be spoken to directly and advised on the symptoms to observe for. Advise they take their child to A&E for further assessment. Written head injury instructions following an injury are good practice, and if available, should be given to the parent (see separate letter) Parents should also be aware that symptoms can be delayed for up to 48 hours post injury.

Most people with concussion make a full and quick recovery. Post-concussion symptoms after a minor brain injury vary; around 20 – 50% have symptoms persisting beyond three months.

Full physical and cognitive rest is important during the recovery period with avoidance of all sport; this also includes swimming and cycling. Cognitive rest includes a rest from watching television, computer games and mobile phone use.

Gradual return to play

The International Rugby Board recommend children and adolescents should not return to play or undertake contact training for a minimum of two weeks following cessation of symptoms. Gradual return to play is a progressive exercise programme, introducing a player back into sport in a controlled and staged manner. It can only be started once the player is symptom free and no longer taking pain- relief for symptom control.

Recommendation

For all PE staff to complete the IRB Concussion Management online resource.

References

1. NHS Choices > Health A-Z > Concussion.

<http://www.nhs.uk/Conditions/Concussion/Pages/Introduction.aspx>

2. Consensus Statement on Concussion in Sport: the 3rd International Conference on Concussion in Sport held in Zurich, November 2008.

3. British Journal of Sports Medicine 2009; 43: i76-i84

4. International Rugby Board > IRB Player Welfare > IRB Concussion Guidelines.

www.irbplayerwelfare **Useful**

websites

www.headway.org.uk – Headway – the brain injury association

Written head injury instructions to be handed to parents following an injury to their child in sport.

Head Injury sustained in sport

Your child sustained a head injury in sport today. If they experience any of the following symptoms, please contact your GP or nearest Accident & Emergency Dept.

Symptoms can be delayed for up to 48 hours post injury.

Increasing headache

visual problems

Increasing drowsiness

Confusion or change in behaviour

Vomiting

Neck stiffness

Fits or convulsions

Consult www.nhs.uk/conditions/head-injury-minor for further information.

Please ask them to attend the medical room when next in school, to ensure they are fit enough to continue with academic and sporting activities. Any concerns, please do not hesitate to contact me.

Mrs Sarah Cadwallader

School Nurse

Telephone - 024 7627 1199

E-mail – nurse@khviii.net

EATING DISORDERS POLICY

Introduction

School staff can play a significant role in recognising and detecting children with early signs of an eating disorder. They also provide support for those affected by this illness. This document describes the approach that King Henry VIII School will take in regards to eating disorders and is intended as guidance for all staff including non-teaching and Governors. **Aims**

- To raise/ increase understanding and promote awareness of eating disorders.
- To help staff recognise the early warning signs of an eating disorder and the leading risk factors.
- To provide staff with support when involved in the management of a pupil suffering with an eating disorder.
- To provide support to pupils suffering or recovering from an eating disorder and their parents/carers and peers.

Definition of eating disorders

Anyone can get an eating disorder regardless of age, sex, or cultural background. It is a complex combination of long standing behavioural, emotional, psychological, interpersonal, and social factors. Food and the control of food are used to compensate for feelings & emotions that may otherwise seem overwhelming. It can become a chronic condition with long term negative effect on relationships, employment, fertility, and parenting.

Risk factors for an eating disorder

Whether or not a person develops an eating disorder will depend on their individual vulnerability, together with the presence of biological or other predisposing factors. **Individual factors**

- Experiencing low self-esteem and other mental health and emotional problems.
- Perfectionist tendencies.
- Adverse sexual experiences and abuse.
- Elevated expectations of achievement, both in school and home.

Environmental factors

- Difficulties in school
- Teasing, critical comments from others about eating, body shape or weight.
- Cultural pressures that glorify “thinness” & the “perfect body”.

Interpersonal factors

- Physical and mental health illness of a close family member including eating disorders, depression, or alcohol misuse.
- Bereavement
- Conflict at home. High parental expectation.

Types of eating disorders

Anorexia Nervosa

Anorexia nervosa is a syndrome in which the individual maintains a low weight because of a preoccupation with body weight, construed either as a fear of fatness or pursuit of thinness. It typically appears in early to mid-adolescence and may present with delayed puberty or stunted growth as well as weight loss.

Signs of Anorexia ☐

Intense fear of weight gain.

- Self-induced weight loss through food avoidance.
- Use of appetite suppressants or laxatives.
- Exercising excessively.
- Distorted body image.
- Menstrual periods stopping and delayed puberty.
- Rigid or obsessive behaviour attached to eating, including eating secretly.

Bulimia Nervosa

Bulimia Nervosa is characterised by recurrent episodes of binge eating and secondly by compensatory behaviour (vomiting, purging, fasting, or exercising or a combination of these) to prevent weight gain. Binge eating is accompanied by a feeling of loss of control over eating. Sufferers will then purge (get rid of food by vomiting or using laxatives). Those who suffer from the non-purging type compensate for binges by exercising or fasting.

Signs of Bulimia Nervosa

- Substantial amounts of food disappearing from the cupboards.
- Going to the bathroom or toilet immediately after meals.
- Using laxatives and vomiting to control weight or sometimes other medications/herbal remedies to lose weight.

Eating Disorders not otherwise specified

Some people may suffer from eating disorders that closely resemble anorexia and bulimia, but are considered atypical, these are classed as Eating Disorders not otherwise specified.

Their weight might be just above the threshold for anorexia or she might still be menstruating. Binge eating and purging may occur less frequently than specified for bulimia. They may have concerns with weight and shape, although in some, primary focus is on maintaining strict control over eating.

Warning signs of an eating disorder

Some eating disorders are mild and can be a passing phase. In others it becomes a long term problem and there is a risk of death. If help is received early during an eating disorder, the better the outcome. Here are some of the first signs that there may indicate a problem: **Physical signs** ☐

- Fainting, headaches, and dizziness.
- Swollen face, around salivary glands ☐ Dry hair & skin; hair loss is common. ☐ Downy hair on face & arms
- Hypersensitivity to heat or cold.
- Tooth decay and/or damaged knuckles

Behavioural changes

- Secretive behaviour & avoiding eating in public.
- Missing meals or eating very little.
- Eating rituals; eating alone, cutting food up into tiny pieces ☒ Wearing baggy clothes to hide body shape.
- Weight controlling behaviour; taking of diuretics, laxatives, diet pill, stimulants ☒ Excessive exercising.

Psychological signs

- Lack of confidence and low self esteem
- Withdrawn and isolated from friends
- Setting unrealistic ambitious standards
- Fatigue and poor concentration including difficult with normal activities.
- Distorted perception of body shape
- Change in mood and personality

Roles and responsibilities of staff

There is a thin line between appropriate responsiveness and inappropriate intrusiveness into the personal lives of students and care needs to be taken not to over step this line.

The goal in school is to detect and address problems in their earliest stages where they exist in thinking and attitudes related to self-image, self-esteem, and self-control. The goal in determining the existence of an eating disorder is simply to raise concerns with the appropriate staff.

Any member of staff, who is aware of a pupil engaging in or suspected to be at risk of an eating disorder, should consult one of the designated teachers for safeguarding children. Please refer to the

Child Protection Policy for King Henry VIII School

Once it is established there may be a concern, an appropriate course of action will be taken.

- Contacting parents/carers
- Arranging a referral to CAMHS, with parental consent.

If a student confides in a member of school staff about their own welfare or that of a peer, they must be made aware that it may not be possible to offer complete confidentiality. If you consider a pupil to be at serious risk of causing themselves harm, then confidentiality cannot be kept. It is important not to make promises of confidentiality that cannot be kept, even if a student puts pressure on you to do so.

Students undergoing treatment for an eating disorder or recovering from an eating disorder All cases will be dealt with on an individual basis in regards to schooling whilst they are suffering with an eating disorder. The decision should be made in discussion with the pupil, their parents, school staff and members of the multi-disciplinary team treating the pupil.

Should the pupil be attending a residential trip, then the trip organiser and nominated health care coordinator will be informed, for health & safety reasons.

If the trip is not residential, then the decision as to whether their information needs to be given will be decided on an individual basis and the nature/duration of the trip. If the pupil does not wish for their health condition to be divulged then they will not be allowed to attend the trip. The reintegration of a pupil into school following a period of absence should be handled sensitively with regular information updates from parents, school staff and a member of the multi-disciplinary team.

Further considerations

Any meetings that take place between pupils, their parents/carers or peers should be recorded in writing to include:

- Dates and times
- An action plan
- Concerns raised
- Details of anyone else who has been informed.

This information should be stored in the pupil's medical records/ My Concern marked confidential and only to be opened in the presence of a member of the Senior Management Team or School Nurse.

SELF-INJURY POLICY

Introduction

Recent research indicates that up to one in ten young people in the UK engage in self-harming behaviours, but such statistics may not reflect the scale of the problem.

This document describes the approach that King Henry VIII School will take in regards to Self-harm behaviours and is intended as guidance for all staff including non-teaching and Governors.

Aims

- To increase understanding and awareness of self-injury.
- To alert staff to warning signs and risk factors.
- To provide support to all staff dealing with students who self-injure.
- To provide support to students who self-harm & and their peers and parents/carers.

Definition of self-injury

Self-harm is a coping strategy. It is a behaviour that should alert us to an underlying problem, difficulty, or disorder. It is a deliberate act and a way of obtaining relief from a difficult and otherwise overwhelming situation or emotional state, but only provides temporary relief; it does not deal with the underlying problems.

Physical pain is often easier to deal with than emotional pain, and its existence can awaken real feelings. It can be a response to everyday stresses and can increase in frequency & severity. It can often become habitual, chronic, and repetitive, but not always for the same reason, or by the same method. An individual can self-harm over a few months or years.

Many acts of self-injury are not directly connected to suicidal intent, in fact the purpose in most cases, is to preserve life and can be a “solution” to an overwhelming problem. People who self-injure usually go to significant efforts to hide their scars or injuries. It can be exceedingly difficult for them to talk about their behaviour, as they may feel ashamed. The act can be very private & personal; it cannot be assumed that it is an attention seeking behaviour.

Although self-harm is not intended as a suicidal act, it must be recognised that the emotional distress that leads to self-harm can also lead to suicidal thoughts and actions.

Self-harm behaviour can include, but is not limited to:

- Cutting, scratching, or picking
- Burning or scalding
- Banging or hitting the head or other parts of the body
- Bruising
- Non suicidal overdose
- Scouring or scrubbing the body excessively.
- Swallowing inedible objects
- Inserting dangerous objects under the skin. ☒ Pulling out hair.

Eating disorders, drug and alcohol misuse, and risk-taking behaviour also fall into the wide definition of self-harm.

Risk factors for self-injury tendencies

The risks of self-injury increase rapidly with the onset of adolescence. It is a manifestation of distress and the presence of other problems. There may be pupils who have none of these risk factors and seem outwardly happy & high achieving, but they may turn to self-injury to cope.

Individual

- Low self esteem
- Onset of a more complicated mental health illness: schizophrenia, Bi-polar disorder, personality disorder & OCD
- Poor communication & problem solving skills.

Environmental

- Being bullied or rejected by peers
- Difficulty in making relationships

Interpersonal

- Neglect or physical, sexual, or emotional abuse ☒ Poor parental and dysfunctional relationships ☒ Frequent arguments with parents.
- Depression, self-injury, or suicide in the family ☒ Unreasonable expectations from family.

Groups more at risk

- Adolescent girls are three times more likely to self-injure than boys, although it is less common in Asian girls.
- Friends of self-harmers seem to be at an increased risk from self-injury.
- Young people struggling with sexual orientation.

Warning signs

Staff may become aware of warning signs that indicate a pupil is experiencing difficulties that may lead to thoughts of self-injury or suicide. Early intervention for self-injury in adolescences is crucial for long-term positive outcomes.

These warning signs should always be taken seriously and staff observing any of these warning signs should seek further advice from the school Nurse or Head of Year.

- Wearing long sleeves & trousers, even in warm weather.
- Increased isolation from friends, being socially withdrawn. ☒ Changes in mood – more aggressive or introverted ☒ Fall in academic achievement.
- Reluctant to take part in physical activities or activities that require a change of clothes.
- Expressing feelings of failure, loss of hope.
- Change in fashion style.
- Bandages to wrists/arms.
- Taking personal risks.
- Abusing drugs & alcohol.

Roles and responsibilities of staff

Students may choose to confide in a member of staff if they are concerned about their own welfare, or that of a peer. Encourage the pupil to be open with you and reassure them that they can get the help they need. Self-injury can be hard for a person to discuss. They may be afraid of how others will react. It is best to respond calmly and non-judgementally, so that the person

remains willing to talk about their self-harming behaviour. Maintaining a pupil's trust is important, if they feel they have lost that trust, they may become more insular and potentially, this can lead to increased self-injury behaviour. Endeavour to enable pupils to feel in control by asking what they would like to happen and what they feel they need. Avoid asking a pupil to stop self-injury, you may be removing the only coping mechanism they have.

Students need to be made aware that it may not be possible for staff to offer complete confidentiality. **If you consider a student is at serious risk of harming themselves then confidentiality cannot be kept.** It is important not to make promises of confidentiality that cannot be kept even if a student puts pressure on you to do so.

Any member of staff, who is aware of a pupil engaging in or suspected to be at risk of engaging in self injury, should consult one of the designated teachers for safeguarding children. Please refer to the Child Protection Policy for King Henry VIII School and log on My Concern.

Following the report, the designated teacher will decide on the appropriate course of action. This may include:

- Contacting parents/guardian
- Arranging a referral to CAMHS with parental consent.
- Immediately removing the pupil from lessons if their remaining in class is likely to cause further distress to themselves or their peers.

In a case of an acutely distressed pupil, the immediate safety pupil is paramount, and an adult should always remain with the pupil.

Further considerations

Any meeting with a pupil, their parents or peers regarding self-harm should be recorded in writing.

☑ Dates and times.

- An action plan
- Areas of concern raised.
- Details of anyone else who has been informed.

This information should be stored in the pupil's file marked confidential and only to be opened in the presence of a member of the Senior Management Team or School Nurse.

Procedure for Blood and Body Fluid Spillage

Purpose

To reduce the risk to staff, student and visitors being exposed to potentially harmful body fluids on the King Henry VIII Senior and Preparatory School.

1. To comply with the requirements of the Health and Safety at Work Act 1974, ensuring employees are not put at risk (section 2).
2. To comply with the Control of Substances Hazardous to Health Regulations. (COSHH).

Definition of Body fluids

Body fluids include; Blood, Urine, Faeces, Vomit, Saliva and Sputum

Procedure

When dealing with Blood and Body Fluid Spillage all staff must always wear personal protective clothing (gloves and aprons).

All protective clothing must be disposed of into Clinical Waste Bags. These are available from the Biology Prep Room.

If there is a risk of extensive splashing of blood and body fluids, extra protection must be worn, such as goggles, face visor and footwear that is fluid repellent.

- Mops should not be used to clear up initial bodily fluid spillages.
- All Body Fluids should be cleaned up immediately and effectively.
- Chemicals used for the cleaning of bodily fluid are hazardous especially when they are used in large volumes in confined spaces or mixed with chemicals or urine. A Control of Substance Hazardous to Health (COSHH) risk assessment must be carried out if using these chemicals.
- Ensure adequate ventilation when using chemical cleaning equipment.
- Disposable gloves must be Nitril or Latex. Any wounds should be covered with a waterproof dressing prior to cleaning any bodily fluids to prevent cross infection.
- Yellow Warning signs must be displayed at all times when dealing with wet floors.

Do not attempt to clean up Body Fluid Spillages unless you have been trained to do so, contact the Caretaker on speed dial 0198, who will come and clean the area with the appropriate chemicals.

Spills on the Body

- ☒ If any part of the body comes into contact with splashes of bodily fluid it must be washed immediately with soap and water.

Reference List

1. Asthma UK (2009) **What is Asthma, What Causes Asthma, AsthmaTriggers?** www.asthma.org.uk
2. Beating Eating Disorders, B-eat (2010) **Understanding disorders and how you can help.** www.b_eat.co.uk
3. BNF (2015) British **National Formulary: 70.** RPS, London.
4. Cleaver B, Crawford R, Armstrong VJ (2006) First **Aid Manual, Step by Step Guide for Everyone.** DK. London. Pg 115, 184-186.
5. Department for Education and Skills (2005) **Managing Medicines in Schools.** DfES Publications. London. Pg 6 (p16), pg 17 (p90), Pg26 (p139), Pg 32 (p186).
6. Diabetes UK (2015) **what is Diabetes? Causes and Risk Factors.** www.diabetes.org.uk
7. Epilepsy Action (2013) **what is Epilepsy?** www.epilepsy.org.uk
8. Ewan PW (1998) Anaphylaxis – **A Clinical Review. ABC of Allergies.** British Medical Journal. 316: 1442-1445.
9. **Good Practice Guidelines for Renal Dialysis/transplantation Units – Prevention and control of Blood-Bourne Virus infection – Public Health Laboratory Service on behalf of Department of Health.** (2002) pg 33 8.25 – 8.26
10. Johanne Lickiss (2007) **Blood and Body Fluid Spillage.** Blackpool, Fylde and Wyre Hospitals Nhs Trust.
11. Kings College London (2010) **Guidelines on Eating Disorders.** www.eatingresearch.com
12. Lee JM, Greenes DS (2000) **BiPhasic Anaphylactic Reactions in Paediatrics.** Paediatrics 106; (4): 762-766
13. National Society for Epilepsy (2013) **About Epilepsy.** www.epilepsysociety.org.uk
14. Newman L, Crawford R, Patterson JG (2001) **First Aid Manual, Emergency Procedures for Everone at Home, Work or at Leisure.** DK, London. Pg 73.
15. NHS Choices (2014) **Asthma.** www.nhs.uk/condition/epilepsy
16. NHS Choices (2013) **Epilepsy.** www.nhs.uk/conditions/epilepsy
17. Patient UK (2015) **Epilepsy – A General Introduction.**

18. Royal College of nursing (2014) **Good practice in Infection Prevention and Control.** Cavendish Square. London.
19. Royal college of Psychiatrists (2008) **Eating Disorders** Royal College of Psychiatrist, London
20. Sampson HA, Mendelson I, Rosen JP (1992) Fatal **and Near Fatal Reactions to Food in Children and Adolescents.** New England Journal of Medicine. 327:380-384.
21. Tall H (2009) **Understanding the causes of disordered eating patterns.** British Journal of School Nursing Vol 4 No 10 pg 487-493
22. The Anaphylaxis Campaign (2015) **Managing Severely Allergic Pupils in School.** The Anaphylaxis Campaign, Hampshire.
23. Tierney LM, McPhee SJ, Papadakis MA (2002). **Current Medical Diagnosis and Treatment.** (int Ed) Lange Medical Books/McCraw Hill New York pg 1203-15.
24. Wilson J (2006) 3rd Edition. **Infection Control in Clinical Practice.** Bailliere Tindall Elsevier.
25. Department of Health March 2015. **Guidance on the use of emergency salbutamol inhalers in School.**
26. Department of Health September 2017 **Guidance on using emergency adrenaline autoinjectors in schools.**
27. **Misuse of Drugs safe custody** (1973 , amendments 2010)
28. **Drugs and Alcohol Education and Prevention** –Ritalin in schools July 2005

APPENDIX A

BKHS King Henry Senior and Preparatory Schools



HOMELY REMEDY PROTOCOL

**HOMELY REMEDIES THAT MAY BE ADMINISTERED TO
STUDENTS OR MEMBERS OF STAFF AT
BKHS KING HENRY SENIOR AND PREPARATORY
SCHOOLS.**

NAME OF MEDICATION	DOSAGE
Paracetamol	120mg – 1g
Ibuprofen	100mg-400mg
Chlorphenamine Maleate (piriton) Loratadine / Certirizine (daily antihistimines)	2mg-4mg
Anthisan (bite/Sting cream)	Sparingly 2-3 times daily
Throat Lozengers sugar free (Strepsils)	No more than 12 daily

HOMELY REMEDY PROTOCOL – PARACETAMOL

Homely Remedy Protocol for administration of **PARACETAMOL** to students or members of staff at King Henry VIII Senior and Preparatory Schools by qualified nurses.

1. Clinical Condition or Situation

Definition of Clinical Situation	<input type="checkbox"/> Management of mild to moderate pain or Pyrexia, as assessed using Nurse's clinical judgement for students or staff at King Henry VIII Senior and Preparatory School.
	<input type="checkbox"/> To include headache, toothache, musculoskeletal pain or other condition warranting simple analgesia.
Persons included	<input type="checkbox"/> Students/staff with mild to moderate pain.
	<input type="checkbox"/> Students/staff with pyrexia above 37.5 °C.
Persons excluded	<input type="checkbox"/> Students/staff to whom paracetamol has been administered within the previous 4 hours.
	<input type="checkbox"/> Students/staff who have taken 4 or more doses of any medication containing paracetamol within the previous 24 hour period.
	<input type="checkbox"/> Students/staff with current hepatic impairment.
	<input type="checkbox"/> Students/staff with current renal impairment.
	<input type="checkbox"/> Students/staff with alcohol dependence.
	<input type="checkbox"/> Students/staff who have had a recent overdose (within the previous 2 weeks) of any substance containing paracetamol.
	<input type="checkbox"/> Students/staff with hypersensitivity to any ingredients of the preparation.
	<input type="checkbox"/> Students/staff on an existing analgesic regime.
Action to be taken in the case of excluded person	Contact parents or GP

2. Staff Competencies	
Authorised staff	<input type="checkbox"/> Registered Nurse Level 1 or authorised consent from the school nurse
Additional requirements	<input type="checkbox"/> Satisfactory completion of Homely Remedy Protocol
	<input type="checkbox"/> To be aware of the medicines available which contain paracetamol
3. Description of Treatment	
Medicines to be administered	Paracetamol 500mg tablets; 500mg soluble tablets, 250mg/5ml suspension, 120mg/5ml.
	Legal status: Tablets = GSL
	Suspension = GSL
	Storage: Locked medical cabinet or trolley
Specific Administration	<input type="checkbox"/> Single dose of 500mg or 1g for adults or children over 16 years.
	<input type="checkbox"/> Single dose of 500mg -750mg for children aged between 12 and 16 years of age.

	<input type="checkbox"/> Single dose of 500mg for children aged between 10 and 12 years of age. <input type="checkbox"/> Single dose of 375mg (7.5mls) for children aged between 8 and 10 years of age. <input type="checkbox"/> Single dose of 250mg (5ml) for children aged between 6 and 8 year of age <input type="checkbox"/> Single dose of 120mg – 240mg for pupils aged between 1 – 5 years of age. Frequency: Every 4-6 hours. Max dose in 24 hours: 4 doses, up to a max of 4g in 24 hours for over 16year olds Max dose of 3mg in 24 hours for children aged 12- 16 years. Max of 2g in 24 hours for children aged 10-12 years. Max of 1g in 24hours for children aged 6 – 10 years.
Follow up treatment	<input type="checkbox"/> Continue to monitor and review until resolved. <input type="checkbox"/> The nurse should use professional judgement to decide when to seek medical intervention.
Warning/Adverse Reactions	<input type="checkbox"/> Side effects rare – rash, blood disorders, liver damage in overdose.
Consent	<input type="checkbox"/> Staff - verbal <input type="checkbox"/> Years 12 and 13 – verbal <input type="checkbox"/> Reception – Year 11 – Parental consent via Health Information Sheet followed by confirmation note sent home advising of administration.
Advice	<input type="checkbox"/> Inform student/member of staff that the medicine is being administered under Homely Remedy Protocol. <input type="checkbox"/> Instruct student/member of staff to avoid other medication containing paracetamol for 4-6 hours.
Record Keeping	<input type="checkbox"/> A record of student’s attendance should be recorded in the medical register.
	<input type="checkbox"/> Record in individual staff/student medical notes, stating date, drug, dosage, time, route. Sign and state administered under Homely Remedy Protocol.

1. Clinical Condition or Situation

Definition of Clinical Situation

- Management of mild to moderate pain or Pyrexia, as assessed using Nurse’s clinical judgement for students or staff at King Henry VIII Senior and Preparatory School.
- To include headache, muscular skeletal pain or other conditions warranting simple analgesia.

Persons included

- Staff/students experiencing mild to moderate pain.

Persons excluded	<input checked="" type="checkbox"/> Staff/students to whom Ibuprofen has been administered within the last 6 hours. <ul style="list-style-type: none"> • Staff/students who have already received 3 or more doses of Ibuprofen within the previous 24 hours. • Any person who is pregnant • Any person who suffers from asthma • Any person with hypersensitivity to aspirin/ibuprofen or other NSAIDS. • Any person with current or previous history of dyspepsia or peptic ulceration. • Any person taking oral anti-coagulants. • Any person suffering from severe cardiac disease. • Any person taking lithium, methotrexate, tacrolimus. • Any person suffering from odema. • Any person suffering from hypertension. • Any person with renal impairment.
Action to be taken in the case of exclusion	<input checked="" type="checkbox"/> Alternative analgesia or advice to seek medical attention.
2. Staff Competencies	
Authorised staff	<input checked="" type="checkbox"/> Registered Nurse Level 1 or with authorised consent from the school nurse Additional Requirements <input checked="" type="checkbox"/> Satisfactory completion of Homely Remedy Protocol Questionnaire.
	<ul style="list-style-type: none"> • To be aware of other medicines that contains Ibuprofen or NSAIDS.
Medicines to be administered	Ibuprofen 200mg tablets, Ibuprofen suspension 100mg/5ml. Legal status: Tablets = GSL Suspension = GSL
Specific Administration	Storage: Locked medicine cabinet or trolley. <input checked="" type="checkbox"/> Single dose of 200-400mg for adults 16 or over. <ul style="list-style-type: none"> • Frequency every 8 hours
Follow up treatment	<input checked="" type="checkbox"/> Max dose in 24 hours: 1.2g <input checked="" type="checkbox"/> Continue to monitor and review until resolved. <ul style="list-style-type: none"> • The nurse should use professional judgement to decide when to seek medical intervention.
Warning/Adverse	<input checked="" type="checkbox"/> Take with or after food.
Reactions	<input checked="" type="checkbox"/> Side effects – rash, abdominal pain, heartburn, diarrhoea, constipation.
Consent	<input checked="" type="checkbox"/> Staff - verbal
Advice	<input checked="" type="checkbox"/> Inform student/member of staff that the medicine is being administered under Homely Remedy Protocol. <ul style="list-style-type: none"> • Instruct student/member of staff to avoid other medication containing ibuprofen for 4-6 hours.
Record Keeping	<input checked="" type="checkbox"/> Record in Ibuprofen Stock Record <ul style="list-style-type: none"> • Record in individual staff/student medical notes, stating date, drug, dosage, time, route. Sign and state administered under Homely Remedy Protocol.

Homely Remedy protocol for Piriton

1. Clinical Condition or situation	
Definition of Clinical Situation	<input type="checkbox"/> Symptomatic relief of allergy, as assessed using nurse’s clinical judgement for students or staff at King Henry VIII Senior and Preparatory School.
	<input type="checkbox"/> For acute allergic rhinitis, acute urticaria and emergency treatment of anaphylactic reactions.
Persons included	<input type="checkbox"/> Staff/students experiencing moderate allergic reactions.
Persons excluded	<input type="checkbox"/> Staff/students with infective rhinitis, renal impairment and hepatic impairment.
	<input type="checkbox"/> Staff/students with hypersensitivity to any ingredients of the preparation.
	<input type="checkbox"/> Persons taking monoamine oxidase inhibitors for depression.
	<input type="checkbox"/> Any person who suffers with epilepsy
	<input type="checkbox"/> Any person who suffers from glaucoma
	<input type="checkbox"/> Any person taking medicines to treat anxiety or to help aid sleep.
	<input type="checkbox"/> Any person suffering from severe cardiac disease
Action to be taken in the case of exclusion	<input type="checkbox"/> Contact parents or emergency services
2. Staff competencies	
Authorised staff	<input type="checkbox"/> Registered Nurse level 1 or with authorised consent from the school nurse
Additional requirements	<input type="checkbox"/> Satisfactory completion of Homely Remedy Protocol Questionnaire
	<input type="checkbox"/> To be aware of other medicines that contains antihistamines.
3. Description of Treatment	
Definition of clinical situation	<p>Piriton 4mg tablets, Piriton suspension 2mg/5ml</p> <p>Legal status: Tablets = P Suspension = P</p> <p>Storage: Locked medicine cabinet or trolley</p>
Specific Administration	<input type="checkbox"/> Single dose of 2mg-4mg for ages 12 and over
	<input type="checkbox"/> Frequency every 4-6 hours
	<input type="checkbox"/> Max dose in 24hours: 24mg
	<input type="checkbox"/> Single dose of 2mg for ages 6-12 years
	<input type="checkbox"/> Frequency every 4-6 hours
	<input type="checkbox"/> Max dose in 24 hours: 12mg
	<input type="checkbox"/> Single dose of 1mg for ages 2-6 years
	<input type="checkbox"/> Frequency every 4-6 hours
Follow up treatment	<input type="checkbox"/> Max dose in 24 hours: 6mg
	<input type="checkbox"/> Continue to monitor and review until resolved <input type="checkbox"/> The nurse should use professional judgement to decide when to seek medical intervention
Warning/Adverse Reactions	<input type="checkbox"/> May cause drowsiness
	<input type="checkbox"/> Side effects- headache, psychomotor impairment, urinary retention, dry mouth, blurred vision and gastro-intestinal disturbances.

	Consent	<input checked="" type="checkbox"/> Staff - verbal
		<input checked="" type="checkbox"/> Years 12 and 13 - verbal
		<input checked="" type="checkbox"/> Years 7 to 11 – Parental consent via Health Information Sheet followed by confirmation sent home advising of administration.
		<input checked="" type="checkbox"/> Nursery to Year 6 only to be used in emergency treatment of anaphylactic reactions.
		<input checked="" type="checkbox"/> Inform student/member of staff that the medicine is being administered under the Homely Remedy Protocol
		<input checked="" type="checkbox"/> Instruct student/member of staff to avoid other medicines containing antihistamines for 4-6 hours

HOMELY REMEDY PROTOCOL- THROAT LOZENGERS

Homely Remedy Protocol for administration of Throat Lozenges to Students or members of staff at King Henry VIII Senior and Preparatory Schools by qualified nurses.

1. Clinical Condition or situation		
	Definition of Clinical Situation	Symptom relief of sore mouth and Throat infections.
	Persons included	Students / Staff with mild to moderate throat discomfort.
	Persons excluded	Student under 6 yrs of age.
		Students / Staff known Diabetic.
		Students /Staff with hypersensitivity to any ingredient within the preparation.
2. Staff competencies		
	Authorised staff	<input checked="" type="checkbox"/> Registered Nurse level 1 or with authorised consent from the school nurse
	Additional requirements	<input checked="" type="checkbox"/> Satisfactory completion of Homely Remedy Protocol
3. Description of Treatment		
	Medicines to be administered	Sugar Free Blackcurrant and menthol lozengers, Strepsils lozengers.
	Specific Administration	Suitable for children 6 years and over. Dissolve 1 lozenge slowly in mouth every 2-3 hours. No more than 12 lozengers in 24 hours.
	Advice	Inform Student / Member of staff that medication is being administered under homely remedy policy.
	Record Keeping	A record of Student's attendance should be recorded in the medical register
		Record in individual Staff / Student medical notes, stating Date, time, drug, dosage, route. Sign and state administered

HOMELY REMEDY PROTOCOL- Anthisan Cream (Bite or Sting relief)

Homely Remedy Protocol for administration of Anthisan Cream to students or members of staff at King Henry VIII Senior and Preparatory School by qualified Nurses.

1. Clinical Condition or situation	
Definition of Clinical Situation	Provide relief from insect bites, stings, nettle rash.
Persons included	Students / Staff with insect bites, stings, nettle rash.
Persons excluded	Students / Staff with signs of skin sensitivity.
	Students / Staff with hypersensitivity to any ingredient within the preparation
2. Staff competencies	
Authorised staff	<input checked="" type="checkbox"/> Registered Nurse level 1 or with authorised consent from the school nurse
Additional requirements	<input checked="" type="checkbox"/> Satisfactory completion of Homely Remedy Protocol
3. Description of Treatment	
Medicines to be administered	Anthisan Cream, Bite and Sting Relief
Specific Administration	Cream should be applied directly to the site of the insect bite, Stinging nettle rash. For best results apply as soon as possible following bite or sting. Apply two to three times daily for up to three days. Do not use of larger areas of skin, Eczema, broken or sunburnt skin. External use only.
Advice	Inform Student / Member of staff that medication is being administered under homely remedy policy.
Record Keeping	A record of Student's attendance should be recorded in the medical register
	Record in individual Staff / Student medical notes, stating Date, time, drug, dosage, route. Sign and state administered

Homely Remedy Protocol - Questionnaire

Name of Nurse.....

Date Completed.....

- To be answered by all Nurses / Deputy Heads wishing to use the protocols.
- The writer to be satisfied with the Nurse / Deputy Head answers prior to being able to implement the protocol.

1. Which **three** medicines can be administered to students/staff under the HRP? Give the name of drug, indication, dose range and frequency.

Drug Name	Indication	Dose Range	Frequency


2. What follow up action would you take after administering a medication under the HRP?

3. What advice would you give to staff/student?

4. Where and what would you document?

5. Under what circumstances would you not administer medication to staff/student under the protocol?

6. What would you do if you administered the wrong dosage of a drug?



APPENDIX C

Instructions for using an auto injector



1. Remove the auto injector from its carry case
2. Grasp the Auto-injector with the tip pointing downwards
3. Pull off grey cap.



4. Place black tip against mid-outer thigh and press firmly until the auto injector activates. Hold in place for several seconds and then remove.
5. Massage the injected area for 10 seconds.



6. Call for an ambulance after administration.



7. Give any used auto-injectors to the Ambulance crew.

DETAILS OF PUPIL

The school will not give your child medicine unless you complete and sign this form, and the Head has agreed that school staff can administer the medication.

Surname

Forename(s).....

Address

.....

Date of Birth:

Class/Form:

Condition or illness:

MEDICATION

Name/Type of Medication (as described on the container)

For how long will your child take this medication:

Date dispensed:

Full Directions for use:

Dosage and method:

Timing:

Special Precautions:

Side Effects:

Self Administration:

Procedures to take in an Emergency:

CONTACT DETAILS:

Name: Daytime Telephone No:

Relationship to Pupil:

Address:

Signed.....Dated.....

(Parents)

Signed.....Dated.....

(Nurse)

Care plan for Temporarily Disabled Pupil/ Staff Member

Pupil/Staff Name	
Form	
Description of Disability	
Aim of Care	Pupil/ Staff Member is able to participate in school activities within reasonable expectation, comfortably and safely

Safety	Measures to be taken in accordance with Temporarily disabled pupil Risk Assessment Form.
Comfort	Wear suitable, comfortable clothing, non-slip shoes. do not apply padding to handles
Personal Care	Self-caring,
Accessibility	Mobilising with crutches. Use accessible entrances/ramps wherever possible. Caution on steps /stairs Organise help to carry belongings etc. Be aware of emergency evacuation procedures/ exits.
Wound Care	

Date Plan initiated	
Initiated by	
Copy to Medical Records Held	
Date Plan updated	
Updated by	
Copy to	